



We create chemistry

ml
800

700

600

500

400

300

200
ml

Basagran® 480 SL

Aqueous Solution
HERBICIDE (KIUAGUGU)

GUARANTEE (DHAMANA):
Bentazone 480 g/l

COMMERCIAL AND AGRICULTURAL CLASS
(KUNDI LA KILIMO NA BIASHARA)

| | | |
|------------------|----|-------------------------|
| GROUP (KUNDI) | C3 | HERBICIDE (KIUAGUGU) |
|------------------|----|-------------------------|

A post-emergence Herbicide for the control of Broad leaf weeds in dry beans, maize and potato.

(Kiuagugu kinachotumika kuzuia magugu yenye matawi mapana kwenye maharwe, vazi mviringo na mahindi.)

Shelf life: Two years from the date of manufacture in tightly sealed original unopened container under cool and dry storage conditions.

(Maisha Rafuni: Miaka miwili kutoka tarehe ya kutengenezwa ikiwa katika chombo chake maalum isipofunguliwa na ikiwe kwa katika pahali pakavu pasipo na joto jingi.)

In case of poisoning call Toll Free number 0800720021/0800730030 (24Hrs).
(Wakati wa madhara piga nambari bila malipo 0800720021/0800730030
(Masaa 24).)

1 L

Net Content (Kipimo Kamili):

WHO Class II: Moderately Hazardous



HARMFUL
(HUDHURU)

READ THE LABEL BEFORE USING
(SOMA KIBANDIKO CHA MAELEZO
KABLA YA KUTUMIA)

KEEP LOCKED OUT OF REACH OF
CHILDREN
(WEKA MBALI NA WATOTO)

REGISTRATION NO. (NAMBARI YA USAJILI): PCPB (CR) 0658

REGISTRANT / MANUFACTURER (MSAJILI / MTENGENEZAJI):
BASF SE, GERMANY

AGENT (AJENTI):

BASF EAST AFRICA LIMITED
P.O. BOX 24271-00100
NAIROBI
TELEPHONE: +254 20 407-2000
Email: info.eastafrica@basf.com

MANUFACTURING DATE: SEE ANOTHER PLACE ON THE LABEL OR PACKAGING
(TAREHE YA KUTENGENEZWA): (MAELEZO YAPO KWINGINE KWENYE
KIBANDIKO HIKI)

BATCH NUMBER: SEE ANOTHER PLACE ON THE LABEL OR PACKAGING
(NAMBARI YA FURUSHI): (MAELEZO YAPO KWINGINE KWENYE KIBANDIKO HIKI)

EXPIRY DATE: SEE ANOTHER PLACE ON THE LABEL OR PACKAGING
(TUMIA KABLA YA): (MAELEZO YAPO KWINGINE KWENYE KIBANDIKO HIKI)



Protect to Provide
Membership Number:
Nambari ya Uanachama: AAK/220



4 014348 931953

81143246 KE 2050

® = Registered trademark of BASF (Alama ya biashara iliyosajiliwa ya BASF)

Basagran® 480 SL

DIRECTIONS FOR USE:

Basagran® 480 SL is a contact post-emergence herbicide for the control of broadleaf weeds in dry beans, maize and potatoes.

MODE OF ACTION

The active ingredient is taken up through the green parts of the plant. The leaves and stems must therefore be adequately wetted with the spray solution. Warmth and good growing weather accelerate the action of the product. Basagran® 480 SL is sprayed post-emergence, i.e. when the weeds have already emerged. After treatment no rain should fall for six hours, so that the active ingredient can penetrate the weeds. Basagran® 480 SL has no biologically active vapour pressure and so constitutes no risk to neighbouring crops in this way.

MIXING INSTRUCTIONS:

Fill the spray tank halfway with water. Start agitator mechanism and slowly add the required amount of Basagran® 480 SL. Add water to the final volume required with continuous agitation.

When tank mixes are to be used, each product should be added separately to the spray tank: Solids first, then Basagran® 480 SL and other liquid herbicides. Use 200-600 l of water. Use the mixture immediately after preparation.

Calibrate the equipment before use and check the nozzle output. During spraying keep the spray solution in motion by means of the agitator or by reflux. After breaks the spray solution must be stirred up again. Over dosage should be avoided. Usual water volume: 200-600 L/ha.

Aerial application is possible with volumes as low as 50 L/ha provided adapted nozzles and pressure produce a droplet spectrum that penetrates adequately into the crop and weeds and is not prone to drift.

RATE OF APPLICATION/SPRAY INTERVAL

| Crop | Target Weed | Rate (L/Ha) | Remarks |
|-----------|--------------------------------|--|--|
| Dry Beans | please see weed spectrum below | 1.5-3.0 L/ha 150-300 ml/20 L Ensure at least 10 Knapsacks/Ha | Treat after formation of the 2nd trifoliate leaf, but not after the flower buds become visible. Slight burning of the foliage may occur, but will grow out again after a short time. There may be non-tolerant varieties. Where a variety has not been tested, it is advisable to carry out a tolerance test before treatment. Apply Basagran® 480 SL in early post-emergence when weeds are young and actively growing. Application should be done when most of the weeds are at 2-4 leaves stage. |
| Potatoes | please see weed spectrum below | 1.75-3.0 L/ha 175-300 ml/20L of water. Ensure a total of 10 Knapsacks/ Ha or 200L water/Ha. | Spray when the weeds are grown and the potatoes are at a height of 10-15 centimeters. Before spraying on large scale, spray a small portion to test for crop tolerance. Apply Basagran® 480 SL in early post-emergence when weeds are young and actively growing. Application should be done when most of the weeds are at 2-4 leaves stage. |
| Maize | please see weed spectrum below | 2.0-3.0 L/ha 200-300 ml/20L of water | Apply Basagran® 480 in early post-emergence when weeds are young and actively growing. Application can be done up to 30 days after crop emergence. |

Weed spectrum

Basagran® 480 SL is effective against the following weeds:

Abutilon theophrasti *Galium aparine*
Acanthospermum hispidum *Helianthus annuus*
Ambrosia artemisiifolia *Heliotropium spp.*
Ambrosia elatior *Hibiscus trionum*
Anoda cristata *Ipomoea purpurea (up to 3-leaf stage)*
Anthemis arvensis *Ipomoea spp.*
Artemisia vulgaris *Kochia scoparia*
Atriplex spp. *Lamium spp. (Up to 3-leaf stage)*
Bidens pilosa *Lapsana communis*
Brassica spp. *Matricaria spp.*
Capsella bursa-pastoris *Mollugo verticillata*
Centaurea cyanus *Phytalis spp.*
Chenopodium album *Polygonum convolvulus*
Chrysanthemum spp. *Polygonum pensylvanicum*
Cyperus esculentus *Richardia spp.*
Datura stramonium *Sida spinosa*
Datura ferox *Sinapis arvensis*
Eclipta spp. *Stellaria media*
Emilia sonchifolia *Tagetes minuta*
Erysimum cheiranthoides *Thlaspi arvense*
Fumaria officinalis *Tribulus terrestris*
Galeopsis tetrahit (up to 3-leaf stage) *Xanthium spp.*
Galinsoga spp.

In the case of root-propagated weeds, only the parts growing above ground are usually controlled. For control of grasses use selective graminicides.

Pre-Harvest-Interval: Beans, Potato – 42 days, Maize – 68 days

Re-Entry-Interval: 24 hours, unless wearing protective clothing.

RESISTANCE MANAGEMENT:

Basagran® 480 SL is inhibitor of photosynthesis on photosystem II (HRAC group C3). As a resistance management practice, this product and products with the same mode of action should be applied in combination or sequence with a herbicide having a different mode of action.

SAFETY PRECAUTIONS:

Ensure thorough ventilation of stores and work areas. When using do not eat, drink or smoke. Hands and/or face should be washed before breaks and at the end of the shift. No special measures necessary if stored and handled correctly.

STORAGE:

Keep away from food, drink and animal feeds. Keep out of the reach of children. Keep away from heat. Protect from direct sunlight.

WASTE DISPOSAL:

On emptying the container, triple rinse thoroughly by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling. Perforate or puncture holes on empty containers and dispose off in accordance with National legislation and Regulation. DO NOT RE-USE empty containers for any other purpose.

LEAK AND SPILLS:

Do not breathe vapour/spray. Use personal protective clothing. Avoid contact with the skin, eyes and clothing. For small amounts: Pick up with suitable absorbent material (e.g. sand, sawdust). For large amounts: Dike spillage. Pump off product. Dispose of absorbed material in accordance with regulations. Collect waste in suitable containers, which can be labeled and sealed. Clean contaminated floors and objects thoroughly with water and detergents, observing environmental regulations. Cleaning operations should be carried out only while wearing breathing apparatus.

ENVIRONMENTAL HAZARDS:

Do not discharge into drains/surface waters/ground water. Do not discharge into the subsoil/soil. Harmful to aquatic life.

FIRST AID MEASURES:

General advice: First aid personnel should pay attention to their own safety. If the patient is likely to become unconscious, place and transport in stable sideways position (recovery position). Immediately remove contaminated clothing.

If Inhaled: Keep patient calm, remove to fresh air and seek medical attention.

Skin contact: After contact with skin, wash immediately with plenty of water and soap. If irritation develops, seek medical attention.

Eye contact: Immediately wash affected eyes for at least 15 minutes under running water, with eyelids open. Consult an eye specialist.

Ingestion: Do not induce vomiting unless told to by a poison control center or doctor. Immediately rinse mouth and then drink 200-300 ml of water, seek medical attention.

TAKE PATIENT TO DOCTOR.

TOXICOLOGICAL INFORMATION:

Signs and symptoms of Poisoning: Causes serious eye irritation. May cause an allergic skin reaction. Harmful if swallowed.

Antidote: Treat according to symptoms (decontamination, vital functions), no known specific antidote.

Note to physician: Treat according to symptoms (decontamination, vital functions).

In case of poisoning call Toll Free number 0800720021/0800730030 (24Hrs).

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use or store a pest control product under unsafe conditions.

WARRANTY:

Seller's guarantee is limited to the terms set out on the label and subject there, to the buyer who assumes the risks to persons or property arising from the use or handling of this product and accepts the product on that condition.

Basagran® 480 SL

MAPENDEKEZO YA UTUMIAJI:

Basagran 480 SL ni kiuagugu kinachotumika kuzuia magugu yenyé matawi mapana kwenye maharagwe, viazi mviringo na mahindi.

JINSI INAVYOFANYA KAZI

Dawa ya Basagran® 480 SL huchukuliwa kupitia kwa matawi na shina za mmea. Hivi basi lazima ikinyunyizwa iħakkishe imeshika matawi na shina vizuri. Basagran® 480 SL yafaa kumunyiziwa kwenye magugu yaliyotoke. Hakikisha ikinyunyizwa kusineshe kwa muda wa masaa sita ndiyo iwezeshe Basagran® 480 SL kupenya ndani ya magugu vizuri.

MAELEZO YA KUCHANGANYA:

Jaza bomba nusu na maji. Ongezea kiasi cha Basagran® 480 SL kinachohitajika. Jaza kabisa bomba na kuanza kukoroga.

Tumia maji kiasi cha lita 200 – 600 Kwa hektari moja. Dawa inaweza kunyuniyiziwa na ndege kwa kiasi cha lita 50 kwa hektari. Dawa inyuniyiziwa mapema wakati magugu yana majani 2-hadi 4 na yasiwe yamepitisha majani 6.

KIPIMO CHA KUTUMIA/MUDA WA KURUDIA

| Mmea | Kipimo (L/hekta) | Maelezo |
|----------------|--|--|
| Maharagwe | Lita 1.5-3.0 kwa hekta Mililita 150-300 / lita 20 za maji. Tumia bomba ya mgongoni isiyó chini ya 10 kwa hekta | Nyunyizia mmea ukipata matawi mawili lakini kabla ya maua kutoke. Kwa vle kuna aina za maharagwe zinazoweza kudhuriwa na dawa hii hakikisha umejaribu kwa mahala kidogo kabla ya unyunyizia shamba loté. |
| Viazi mviringo | Lita 1.75-3.0 kwa hekta Mililita 60-100 / Lita 20 za maji. Hakikisha umetumia bomba 30 au maji lita 600 kwa Hekta | Nyunyizia magugu yakimea na mmea wa viazi ukifikisha urefu wa centimita 10-15. Kwa vle kuna aina za viazi zinazoweza kudhuriwa na dawa hii, hakikisha umejaribu kwa mahala kidogo kabla ya kunyuniyizia shamba loté. |
| Mahindi | Tumia lita 2-3 kwa hekta Mililita 200-300 / Lita 20 za maji. | Nyunyizia Basagran® 480 SL mapema wakati magugu ni wachanga na wanamea kwa kasi. Unyunyizaji unaweza kufanyika siku thelathini baada ya mazao kuibuka |

Aina ya magugu

| | |
|---|--|
| Basagran® 480 SL kinaweza kuzuia magugu yenyé matawi mapana Kama ifuatavyo: | |
| <i>Abutilon theophrasti</i> | <i>Galium aparine</i> |
| <i>Acanthospermum hispidum</i> | <i>Helianthus annuus</i> |
| <i>Ambrosia artemisiifolia</i> | <i>Heliotropium spp.</i> |
| <i>Ambrosia elatior</i> | <i>Hibiscus trionum</i> |
| <i>Anoda cristata</i> | <i>Ipomoea purpurea (up to 3-leaf stage)</i> |
| <i>Anthemis arvensis</i> | <i>Ipomoea spp.</i> |
| <i>Artemisia vulgaris</i> | <i>Kochia scoparia</i> |
| <i>Atriplex spp.</i> | <i>Lamium spp. (up to 3-leaf stage)</i> |
| <i>Bidens pilosa</i> | <i>Lapsana communis</i> |
| <i>Brassica spp.</i> | <i>Matricaria spp.</i> |
| <i>Capsella bursa-pastoris</i> | <i>Mollugo verticillata</i> |
| <i>Centaurea cyanus</i> | <i>Physalis spp.</i> |
| <i>Chenopodium album</i> | <i>Polygonum convolvulus</i> |
| <i>Chrysanthemum spp.</i> | <i>Polygonum pensylvanicum</i> |
| <i>Cyperus esculentus</i> | <i>Richardia spp.</i> |
| <i>Datura stramonium</i> | <i>Sida spinosa</i> |
| <i>Datura ferox</i> | <i>Sinapis arvensis</i> |
| <i>Eclipta spp.</i> | <i>Stellaria media</i> |
| <i>Emilia sonchifolia</i> | <i>Tagetes minuta</i> |
| <i>Erysimum cheiranthoides</i> | <i>Thlaspi arvense</i> |
| <i>Fumaria officinalis</i> | <i>Tribulus terrestris</i> |
| <i>Galeopsis tetrahit (up to 3-leaf stage)</i> | <i>Xanthium spp.</i> |
| <i>Galinsoga spp.</i> | |

Muda salama wa kungoja kabla ya kuvuna: Mahagwe, viazi-siku 42, Mahindi-siku 68

Muda wa kungoja kabla kuingia mahali paliponyonyiziwa dawa: Masaa 24 ama kama ulo na nguo za kujikinga

KUZUIA UZOEFU WA KUVU:

Basagran 480 SL ina kiambato amalifu ya kikundi cha magugu walio katika Kikundi ya HRAC C3. Kwa kuzuia uzoefu wa magugu kutokana na mchanganyiko huu, viuagugu hivi na vingine yaa kikundi kimba havifai kutumika moja baada ya nyininge kwenye msimu mmoja. Fuatanisha na viuagugu vingine yaa vikundi tofauti.

TAHADHARI KWA MTUMIAJI:

Hakikisha uingizaji wa hewa ni mzuri kwenye sehemu za kuhifadhi dawa na maeneo ya kazi. Usile, usinywe wala usivute sigara unapotumia dawa hii. Osha mikono na uso iliyoquswa, kabla kwenda kwa haja na baada ya kumaliza kutumia dawa. Hakuna hatua maalum zinazohitajika ikiwa zimehifadhiwa na kushughulikiwa kwa usahihi.

KEHIFADHI:

Weka dawa mbali na chakula, vinywaji na mifugo. Weka dawa mbali na mahali panapoweza kutokea moto. Linda dawa dhidi ya jua ya moja kwa moja. Weka dawa mbali na watoto.

KUTUPA MABAKI:

Osha chombo cha dawa mara tatu na umwage ule mwosho kwenye tanki la kunyuyizia. Haribu chombo cha dawa kwa kutoboa mashimo baada ya kuitumia dawa yote iliyo halafu itupe mahali pazuri mbali na mto kuambatana na sheria zilizoko kwenye nchi. USITUMIE KWA MATUMIZI INGINIE.

DAWA IKIMWAGIKA:

Usipumue mvuke/dawa unapotumia dawa hii. Vaa mavazi ya kujikinga kama ulivyeolezwa hapo juu. Unapotumia dawa hii linda macho, ngozi na nguo ulizovaa. Kwa kiasi kidogo: Pangusa dawa iliymwagika na mchanga au yumbi inayopatikana wakati mbaao inapopasuliwa. Kwa kiasi kubwa: Pampu dawa iliymwagika. Tupa vifaa vilivyoonyanza dawa kulingana na sheria zilizoko kwenye nchi. Kusanya taka katika vyombo vinayofaa, ambavyo vinawenza kuwekwa lebo na sili. Safisha sakafu iliymwagika na dawa kwa maji mengi yaliyo na sabuni, kufutana na kanuni za mazingira. Shughuli za kusafisha zinapaswa kufanyika tu wakati wa kuvala vifaa nya kupumua.

HADHARI KWA MAZINGIRA:

Usichafue vidimbwi nya maji kwa dawa hii au mabaki yake. Usioshe vyombo vilivytumika karibu na mito. Usimwage mabaki kwenye udongo au mito. Inaweza leta madhara kwa wanyama wa majini na viumbe vingine wanaoishi kwenye maji.

HUDUMA YA KWANZA:

Mashauri: Watoa huduma ya kwanza wanapaswa kuzingatia usalama wao wenye. Ikiwa mgongajiwa anaweza kukosa fahamu, msafirishie ikiwa amelalia upande (nafasi ya kupona). Ondoa nguo zilizishika dawa mara moja.

Ikiwa imeingizwa: Weka mgongajiwa atulie, mueke kwenye hewa safi na utafute matibabu.

Kushikwa kwa Ngozi: Baada ya kuguswa na dawa kwa ngozi. Osha kwa maji mengi na sabuni. Kuwashwa kwa ngozi kukitokea, patua ushauri wa matibabu.

Kushikwa kwa Macho: Osha macho yaliyoathiriwa kwa muda wa dakika 15 chini ya maji ya kukimbia na kope la jicho likifunguliwa wazi. Wasiliana na mtaalamu wa jicho.

Kumeza: Usiruhusu kutapika isipokuwa kama umeambwa na kituo cha udhibiti wa sumu au daktari. Mara moja suuu mdomo na kisha kunya mililita 200-300 ya maji, tafuta matibabu.

PELEKA MJERUHI KWA DAKTARI.

MAELEZO KUHUSU SUMU:

Dalili ya kusumika: Inasababisha kuwashwa sana kwa jicho. Inasababisha kuwashwa sana kwa ngozi. Inadhuru ikimezwu.

Kiuasumu: Hakuna tiba ijulikanayo kwa sasa. Tibu kufuatia dalili za ugonjwa.

Maagizo kwa daktari: Tibu mgongajiwa kulingana na dalili za ugonjwa.

Wakati wa madhara piga nambari bila malipo 0800720021/0800730030 (Masaa 24).

ILANI KWA WATUMIAJI:

Dawa hii sharti itumiwe kulingana na maagizo kwenye kibandiko hiki tu. Ni hatia chini ya Sheria za madawa ya kuangamiza wadudu waharibifu kutumia au kuweka dawa hii mahala pasipo usalama.

THIBITISHO:

Dhamana ya Muuzaji ni maelezo yaliyo kwenye kibandiko ambayo ni lazima yafuatwe kikamilifu. Mnunuzi lazima akubali hatari zozote zinawenza kutokana na matumizi ya dawa hii kwa binadamu au mali na lazima aikubali dawa hii katika hali hiyo.



We create chemistry

ml
800

700

600

500

400

300

200
ml

Basagran® 480 SL

Aqueous Solution
HERBICIDE (KIUAGUGU)

GUARANTEE (DHAMANA):
Bentazone 480 g/l

COMMERCIAL AND AGRICULTURAL CLASS
(KUNDI LA KILIMO NA BIASHARA)

| | | |
|------------------|----|-------------------------|
| GROUP (KUNDI) | C3 | HERBICIDE (KIUAGUGU) |
|------------------|----|-------------------------|

A post-emergence Herbicide for the control of Broad leaf weeds in dry beans, maize and potato.

(Kiuagugu kinachotumika kuzuia magugu yenyenye matawi mapana kwenye maharawe, viasi mviringo na mahindi.)



Shelf life: Two years from the date of manufacture in tightly sealed original unopened container under cool and dry storage conditions.

(Maisha Rafuni: Miaka miwili kutoka tarehe ya kutengenezwa ikiwa katika chombo chake maalum isipofunguliwa na ikiwe kwa katika pahali pakavu pasipo na joto jingi.)

In case of poisoning call Toll Free number 0800720021/0800730030 (24Hrs).
(Wakati wa madharra piga nambari bila malipo 0800720021/0800730030
(Masaa 24.).

1 L

Net Content (Kipimo Kamili):

WHO Class II: Moderately Hazardous



HARMFUL
(HUDHURU)



READ THE LABEL BEFORE USING
(SOMA KIBANDIKO CHA MAELEZO
KABLA YA KUTUMIA)

KEEP LOCKED OUT OF REACH OF
CHILDREN
(WEKA MBALI NA WATOTO)

REGISTRATION NO. (NAMBARI YA USAJILI): PCPB (CR) 0658

REGISTRANT / MANUFACTURER (MSAJILI / MTENGENEZAJI):
BASF SE, GERMANY

AGENT (AJENTI):

BASF EAST AFRICA LIMITED
P.O. BOX 24271-00100
NAIROBI
TELEPHONE: +254 20 407-2000
Email: info.eastafrica@bASF.com

MANUFACTURING DATE: SEE ANOTHER PLACE ON THE LABEL OR PACKAGING
(TAREHE YA KUTENGENEZWA): (MAELEZO YAPO KWINGINE KWENYE KIBANDIKO HIKI)

BATCH NUMBER: SEE ANOTHER PLACE ON THE LABEL OR PACKAGING
(NAMBARI YA FURUSHI): (MAELEZO YAPO KWINGINE KWENYE KIBANDIKO HIKI)

EXPIRY DATE: SEE ANOTHER PLACE ON THE LABEL OR PACKAGING
(TUMIA KABLA YA): (MAELEZO YAPO KWINGINE KWENYE KIBANDIKO HIKI)



Protect to Provide
Membership Number:
Nambari ya Uanachama: AAK/220



® = Registered trademark of BASF (Alama ya biashara iliyosajiliwa ya BASF)

81143246 KE 2050